

Key Nauru DHS Indicators



	Total
Marriage and fertility	
1. Women aged 20–24 married by age 18 (%)	27
2. Men aged 20–24 married by age 18 (%)	12
3. Total fertility rate (children per woman)	3.4
4. Women aged 15–19 who were already mothers or pregnant at the time of the survey	15
5. Median age at first birth for women aged 25–49	21.9
6. Married women (15–49 years) with 2 living children wanting no more children (%)	24
Family planning (currently married women aged 15 – 49)	
Current use of family planning	
7. Any method (%)	35.6
8. Any modern method	25.1
8a. Female sterilisation	13.3
8b. Male sterilisation	0.2
8c. Injectables	2.3
8d. Pill	0.6
8e. Condom	2.8
Unmet need for family planning	
9. Total unmet need	23.5
9a. For spacing	16.4
9b. For limiting	7.1
Infant and child mortality	
For the 10-year period preceding the survey	
10. Neonatal mortality	27
11. Infant mortality rate	39
12. Under-five mortality rate	41
For the 5-year period preceding the survey	
13. Neonatal mortality	27
14. Infant mortality rate	38
15. Under-five mortality rate	38
Maternal and child health	
Maternity care for births in the last 3 years	
16. Mothers who had at least 4 antenatal care visits for their last birth (%)	40.2
17. Births assisted by a doctor/nurse/other health professional	97.4
18. Births in a hospital or health facility	98.7
19. Mothers who received postnatal care within 2 days of delivery for their last birth	65.6
Child immunisation and vitamin A supplementation	
20a. Children 12–23 months fully immunised (BCG, measles, and 3 doses each of polio/DPT) (%)	85.5
20b. Children 12–23 months who have received BCG (%)	98.4
20c. Children 12–23 months who have received 3 doses of polio vaccine (%)	91.0
20d. Children 12–23 months who have received 3 doses of DPT vaccine (%)	89.0
Treatment of childhood diseases (children under 3 years)	
21. Children with diarrhoea in the last 2 weeks who received ORS	23.3
22. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	34.3
23. Children with fever in the last 2 weeks taken to a health facility (%)	50.6
Child feeding practices and nutritional status of children	
24. Children under 3 years breastfed within 1 hour of birth (%)	76.4
25. Children 0–5 months exclusively breastfed (%)	(67.2)
26. Children under 3 years who are stunted (%)	24
27. Children under 3 years who are wasted (%)	1
28. Children under 3 years who are underweight (%)	4.8
Knowledge of HIV and AIDS among ever married adults (aged 15–49)	
29. Women who have heard of AIDS (%)	73.1
30. Men who have heard of AIDS (%)	83.2
33. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	56.2
34. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	66.7
36. Women who had high risk sex in the past 12 months (%)	23.9
35. Men who had high risk sex in the past 12 months (%)	51.5